

USE OF COMMON CAROTID INTIMA-MEDIA THICKNESS MEASURED BY ULTRASOUND ECHO-TRACKING IN CARDIOVASCULAR RISK STRATIFICATION BEFORE NON-CARDIAC SURGERY IN LOW-RISK CATEGORY: A RESEARCH IDEA

Srinivas Mantha, MD*, Sudha Lakshmi Tripuraneni, MD†‡, Lee A Fleisher, MD‡‡, Michael F. Roizen, MD¶¶_*Consultant Anesthesiologist & Pain Physician, Mantha’s Pain Clinic, Hyderabad, India; † Consultant Cardiologist, Mantha’s Heart Clinic, Hyderabad, India; ‡‡Department of Anesthesiology and Critical Care, University of Pennsylvania, Philadelphia, USA, ¶¶_Chief Wellness Officer, Roizen Family Chair, The Wellness Institute of Cleveland Clinic, Cleveland, Ohio, USA

ABSTRACT

Background: Preoperative cardiac risk stratification algorithms aim to reduce postoperative morbidity and mortality. They typically use a Bayesian approach to identify a low-risk category group who can go for surgery without further testing. A recent study found that three popular prediction models disagreed 29% of the time by which patients were categorized as low risk (<1%) [1]. Hence, an approach to strengthen and optimize the Bayesian risk indices is needed [2]. Common Carotid Intima-Media Thickness (CCIMT) measured by Ultrasound is a surrogate marker for atherosclerosis and quantifies atherosclerotic burden in entire vascular tree. Two variables of particular interest i.e. CCIMT-z score and vascular age help in such quantification [3]. From a clinical series in an outpatient cardiology clinic, information obtained from CCIMT was explored for feasibility in risk stratification.

Methods: As a part of clinical or wellness care, CCIMT was measured by B-mode ultrasonography using 3-13 MHz linear probe. An accurate method called ‘echo-tracking’ that relies on automated edge detection by radiofrequency signal processing of ultrasound was used (Figure). From a series of 44 cases, 22 were segregated who would otherwise qualify for low-risk category should they present for preoperative evaluation for noncardiac surgery. CCIMT z scores and percentiles for vascular age were computed based on population-based normal values at different ages in either gender. A multivariate linear regression analysis was done with CCIMT z score as dependent variable and following as independent variables: body mass index (BMI), waist-to-height ratio (WHR), total cholesterol HDL ratio (TC-HDL ratio) and serum vitamin D3 levels. According to American Society of Echocardiography, a CCIMT z score of ≥1.96 equivalent to ≥97.5 percentile is defined as highly abnormal that requires immediate attention and further evaluation.

Results: Mean (SD) for age were 48 years (12) and there were equal number of men and women. Of the independent variables tested, TC-HDL ratio was significantly associated with increased CCIMT z score (p=0.0134). A simple linear regression analysis using only the TC-HDL ratio yielded the model: IMT_Zscore = -0.1149 + (0.3400 x TC-HDL_Ratio). The p value for the slope was 0.0033 and correlation coefficient was 0.5976 and R square was 0.3571. There were four cases in this low-risk group who had z score ≥1.96

Discussion: CCIMT appears to be useful as part of shared decision making to determine if long-term survival is important with respect to the decision to operate. Preoperative statin therapy and close monitoring with troponin may be considered to reduce immediate perioperative risk. Hence, potential application of CCIMT in cardiac risk stratification appears promising. A suitable model to predict the CCIMT z score using readily available clinical and laboratory variables could be developed in the research setting. In addition, scope of substituting vascular age for chronological age and/or categorizing individuals with Z score ≥1.96 as having severe systemic disease in existing risk stratification algorithms may be explored. We believe that the suggested approach would provide better insights for future research in the area.

- Glance LG, Faden E, Dutton RP, Lustik SJ, Li Y, Eaton MP, Dick AW. Impact of the Choice of Risk Model for Identifying Low-risk Patients Using the 2014 American College of Cardiology/American Heart Association Perioperative Guidelines. Anesthesiology 2018;129:889-900.
- Fleisher LA. Preoperative Cardiac Evaluation before Noncardiac Surgery: Reverend Bayes's Risk Indices and Optimal Variables. Anesthesiology. 2018;129(5):867-8
- Engelen L, Ferreira I, Stehouwer CD, Boutouyrie P, Laurent S, Reference Values for Arterial Measurements C. Reference intervals for common carotid intima-media thickness measured with echotracking: relation with risk factors. Eur Heart J 2013;34:2368-80.




Scanning for CCIMT


- Longitudinal view of CCA
- Tuning Fork Appearance
- Double line sign in far wall of CCA which represent intima and media

INTRODUCTION

Introduction:


- Preoperative cardiac risk stratification algorithms aim to reduce postoperative morbidity and mortality
- They typically use a Bayesian approach to identify a low-risk category group who can go for surgery without further testing
- Separation of *low-risk* (<1%) from *elevated-risk* (≥1%) is the key decision point in the ACC/AHA guidelines for preoperative evaluation for non-cardiac surgery A recent study found that three popular prediction models disagreed 29% of the time by which patients were categorized as low risk (<1%) [1]. The three popular models are
 - The Revised Cardiac Risk Index (Lee et al. Circulation 1999;100:1043-9.
 - The American College of Surgeons National Surgical Quality Improvement Program Surgical Risk Calculator (Bilimoria et al. J Am Coll Surg 2013;217:833
 - Risk calculator for prediction of cardiac risk after surgery. Gupta et al. Circulation 2011;124:381
- Hence, an approach to strengthen and optimize the Bayesian risk indices is needed [2].
- Common Carotid Intima-Media Thickness (CCIMT) measured by ultrasound is a surrogate marker for atherosclerosis and quantifies atherosclerotic burden in entire vascular tree.
- As CCIMT increases with age , absolute values are difficult to interpret in a given individual. Hence, two objective variables of particular interest i.e. CCIMT-z score and vascular age help in such quantification [3].
- From a clinical series in an outpatient cardiology clinic, information obtained from CCIMT was explored for feasibility in risk stratification.





Echo-tracking and auto-edge detection


- Real time radiofrequency processing of ultrasound
- Region of interest (ROI) is 1.5 cm starting from 1 cm of vertical reference line just proximal to carotid bulb
- A table along side of image gives measurements of last 6 cardiac cycles
- Good quality measurement indicators: SD less than 10 and A thick green overlay within ROI
- The method is inherited by Esaote **ART.LAB** experience and now available in all MyLab series of Esaote (Italy) ultrasound machines



The technique enhances Accuracy & Reliability

METHODS and RESULTS

Methods: As a part of clinical or wellness care, CCIMT was measured by B-mode ultrasonography using 3-13 MHz linear probe. An accurate method called ‘echo-tracking’ that relies on automated edge detection by radiofrequency signal processing of ultrasound was used . From a series of 44 cases, 22 were segregated who would otherwise qualify for low-risk category should they present for preoperative evaluation for noncardiac surgery. CCIMT z scores and percentiles for vascular age were computed based on population-based normal values at different ages in either gender. A multivariate linear regression analysis was done with CCIMT z score as dependent variable and following as independent variables: body mass index (BMI), waist-to-height ratio (WHR), total cholesterol HDL ratio (TC-HDL ratio) and serum vitamin D3 level




A CCIMT z score of ≥1.96 equivalent to ≥97.5 percentile is defined as highly abnormal that requires immediate attention and further evaluation.

Results:

- Mean (SD) for age were 48 years (12) and there were equal number of men and women.
- Of the independent variables tested, TC-HDL ratio was significantly associated with increased CCIMT z score (p=0.0134).
- A simple linear regression analysis using only the TC-HDL ratio yielded the model: IMT_Zscore = -0.1149 + (0.3400 x TC-HDL_Ratio).
- The p value for the slope was 0.0033 and correlation coefficient was 0.5976 and R square was 0.3571.
- There were four cases in this low-risk group who had z score ≥1.96

A website designed by the first author (SM), www.suhitam.com/vascularage takes four input variables i.e. age, gender and CCIMT in µm of either sides and yields a clinically useful report . The algorithm uses maximum CCIMT of the two sides. Report of a male individual aged 40 years with CCIMT values of 700 and 730 µm respectively on right and left sides, will be as follows:



IMT- Meas	IMT- Std	Z- score	Perce ntile	Vasc age	Risk Grade
730	532	2.21	98	78	4

Visitors of the website can login using ‘*visitor*’ for both user id and password to view representative cases with different risk grades based on CCIMT.

DISCUSSION and REFERENCES

Discussion

- Potential application of CCIMT in cardiac risk stratification appears promising. A suitable model to predict the CCIMT z score using readily available clinical and laboratory variables could be developed in the research setting. In addition, scope of substituting vascular age for chronological age and/or categorizing individuals with Z score ≥1.96 as having severe systemic disease in existing risk stratification algorithms may be explored. We believe that the suggested approach would provide better insights for future research in the area.
- CCIMT has the potential to be incorporated into shared decision making [4] to provide information on long-term survival to incorporate into the decision to operate.
- CCIMT results can also be used for the decision to implement Preoperative statin therapy and close monitoring with troponin to reduce immediate perioperative risk.
- CCIMT measurement can be performed by anesthesiologist during preoperative evaluation and expand our role as perioperative physicians. If the stated machine is not available, methods such as using electronic calipers on frozen images, other techniques of auto-edge detection available in other machines may be used. Advantages of primary care taker in direct visualization of intima , the abnormality of which is responsible for vascular events, is obvious.
- A prospective study is planned by the authors to identify determinants of CCIMT in healthy individuals. Protocol summary may be viewed from a clinical trial registry using ‘ccimt’ in the key word search <http://ctri.nic.in/Clinicaltrials/advancesearchmain.php>

References

- Glance LG, Faden E, Dutton RP, Lustik SJ, Li Y, Eaton MP, Dick AW. Impact of the Choice of Risk Model for Identifying Low-risk Patients Using the 2014 American College of Cardiology/American Heart Association Perioperative Guidelines. Anesthesiology 2018;129:889-900.
- Fleisher LA. Preoperative Cardiac Evaluation before Noncardiac Surgery: Reverend Bayes's Risk Indices and Optimal Variables. Anesthesiology. 2018;129(5):867-8
- Engelen L, Ferreira I, Stehouwer CD, Boutouyrie P, Laurent S, Reference Values for Arterial Measurements C. Reference intervals for common carotid intima-media thickness measured with echotracking: relation with risk factors. Eur Heart J 2013;34:2368-80.
- Raymond BL, Wanderer JP, Hawkins AT, Geiger TM, Ehrenfeld JM, Stokes JW,McEvoy MD. Use of the American College of Surgeons National Surgical Quality Improvement Program Surgical Risk Calculator During Preoperative Risk Discussion: The Patient Perspective. Anesth Analg 2019;128:643-50.

Author disclosures: Conflicts of Interest: None